

JSNA Health and Wellbeing Profile 2025/26

Cardiovascular Disease (CVD)

Summary points

- In the 3 years period of 2021-2023 there were 818 premature (aged under 75 years) deaths from cardiovascular disease (CVD) in Bristol. 70.4% of these were among men.
- The under 75 mortality rate from cardiovascular diseases in Bristol was 87.1 per 100,000 population in 2021-2023, significantly higher than the England average of 77.1.
- Among those who died of CVD prematurely, 382 died of coronary heart disease (rate 41.2 per 100,000) and 115 died of stroke (rate 12.2 per 100,000).
- CVD premature mortality rates vary significantly across the city. In the Inner City locality, rates have been consistently above the Bristol average and over 3 times as high as the rates in the North and West (inner) locality.
- Early death rates from CVD are increasing.

Cardiovascular Disease mortality

Cardiovascular disease (CVD) is one of the major causes of death in under 75s in England.

Early deaths due to CVD were declining nationally from 2001, but this decline has now levelled off. Since 2020 the rates started to increase. In Bristol rates have levelled off for both males and females, with most recent data showing an increase in rates among males.

In 2021-2023, there were 818 early (under 75) deaths from CVD in Bristol, a rate of 87.1 deaths per 100,000 population. This is statistically significantly higher than the England rate of 77.1 per 100,000¹.

In Bristol, the rate of under 75 deaths from CVD among males at 124.3 per 100,000 is over double the rate among females (50.8 per 100,000). A similar difference is seen nationally.




Figure 1: Under 75 mortality rates from all cardiovascular diseases, [Mortality Profile - Data | Fingertips | Department of Health and Social Care](#), July 2025

¹ Source: [Mortality Profile - Data | Fingertips | Department of Health and Social Care](#), July 2025

Local data² on variation across the city shows the North & West (inner) locality rates are significantly lower than the Bristol average, and the rates are highest among males in the Inner City (Figure 2).




Figure 2: Under 75 mortality rates from all cardiovascular diseases, Primary Care Mortality Database via NHS England

Between 2011-2013 and 2018-2020 the rates of under 75 mortality from CVD have fallen in all localities in Bristol. However, from the 2019-2021 period the rates have been increasing in all localities. The rates in the Inner City locality have been consistently above the Bristol average and over 3 times as high as the rates in the North and West (inner) locality (Figure 3).




Figure 3: Under 75 mortality rates from all cardiovascular diseases, Primary Care Mortality Database via NHS Digital

² 2021-2023 locality data from the Primary Care Mortality Database via NHS England

Coronary Heart Disease (CHD)

Data from GP registers³ show that recorded prevalence of Coronary Heart Disease (CHD) across Bristol and in England in 2023/24 is similar to the previous year's figures.

Bristol recorded prevalence (2.2%) is lower than the England average (3.0%). However, in the North & West (outer) locality the recorded prevalence of CHD is at England's level of 3.0%. While this is over double the rate of the Inner City locality (Figure 4), these data are not age standardised and the lower recorded prevalence in Inner City may in part reflect the higher proportion of younger people living in Inner City than in other localities.




Figure 4: CHD prevalence, NHS Quality Outcomes Framework 2023/24, via NHS England

Local data for early deaths from CHD show rates in North and West (inner) locality remain significantly lower than the city average, and are less than half the rates of the Inner City and North and West (outer) localities.

Overall CHD early death rates in Bristol are over 3 times higher for men than for women (Figure 5).

Emergency hospital admissions for CHD during 2023/24 record 63.2% of Bristol patients were male and 36.8% were female⁴.

³ Source: NHS England QOF data for 2022/23.

⁴ Hospital Episode Statistics via NHS England




Figure 5: Under 75 mortality rates from coronary heart disease (CHD), Primary Care Mortality Database via NHS England

Stroke

There were 116 early deaths from stroke in 2021-2023 in Bristol. The under 75 death rate from stroke is highest in the Inner City, and lowest in the South locality. At locality level the figures have wide confidence intervals and are statistically similar to Bristol average.

Overall, in Bristol, over 17% more men than women die early from stroke (Figure 6).




Figure 6: Under 75 mortality rates from stroke, Primary Care Mortality Database via NHS England

High Blood Pressure (Hypertension)

Hypertension increases risk of heart disease or stroke. In 2023/24 there were 63,096 patients with hypertension on Bristol's GP registers. Prevalence of recorded hypertension varies across the city, with highest rates in the South and North & West (outer), and lowest in Bristol East and the Inner City (Figure 7). Overall the recorded prevalence of hypertension in Bristol is lower than in England on average. Note these data are not age-standardised.




Figure 7: Hypertension prevalence, NHS Quality Outcomes Framework 2023/24, via NHS England

Equalities:

CVD is one of the conditions most strongly associated with socioeconomic deprivation. Of the most common causes of death, CVD is the largest contributor to the gap in life expectancy between the most and least deprived, accounting for up to 25% of the difference. If you live in England's most deprived areas, you are almost four times as likely to die prematurely than those in the least deprived. CVD is also more common where a person is male, older, has a severe mental illness, or ethnicity is South Asian or African Caribbean.⁵

In Bristol the rate of early deaths from CVD is 2.7 times higher among people living in the most deprived areas of the city compared to the most affluent areas (Figure 8)

⁵ Public health matters: Health Matters: Preventing cardiovascular disease.

<https://publichealthmatters.blog.gov.uk/2019/02/14/health-matters-preventing-cardiovascular-disease/>




Figure 8: Under 75 mortality rates from CVD by deprivation quintile, Bristol 2020-2022. Primary Care Mortality Database via NHS England

Further data / links:

- Public Health Outcomes Framework: <https://fingertips.phe.org.uk/profile/public-health-outcomes-framework>
- Quality Outcomes Framework: [Quality and Outcomes Framework - NHS England Digital](#)
- Public Health England Cardiovascular Disease, Diabetes and Kidney Disease Profiles: <https://fingertips.phe.org.uk/profile-group/cardiovascular-disease-diabetes-kidney-disease>

Covid-19 impact:

There is now clear evidence that Cardiovascular disease (CVD) and COVID-19 share common underlying risk factors. 'People with pre-existing CVD, diabetes, obesity and high blood pressure are not only at greater risk of having a heart attack, stroke or developing some forms of dementia, but are known to experience more severe outcomes from COVID-19, including hospitalisation, ventilation and death.'⁶

Date updated: July 2025

Date of next update: May 2026

⁶ Public Health Matters: Cardiovascular disease: building back better.
[Cardiovascular disease: building back better - Public health matters \(blog.gov.uk\)](#)