


PYROCORE

PHOENIX AND DEMO SMALL WASTE INCINERATION PLANT:

HUMAN HEALTH RISK ASSESSMENT



September 2022

Report Reference: C98-P11-R01

CONTENTS

| | | |
|----------------|--|-----------|
| 1 | INTRODUCTION..... | 1 |
| 1.1 | BACKGROUND..... | 1 |
| 1.2 | DESCRIPTION OF THE INSTALLATION..... | 1 |
| 1.2.1 | <i>Introduction.....</i> | 1 |
| 1.2.2 | <i>Phoenix SWIP.....</i> | 2 |
| 1.2.3 | <i>Demonstrator Unit.....</i> | 2 |
| 1.3 | PURPOSE OF THE ASSESSMENT..... | 3 |
| 1.4 | SCOPE OF THE ASSESSMENT..... | 4 |
| 1.5 | APPROACH TO THE ASSESSMENT..... | 5 |
| 2 | METHODOLOGY FOR ESTIMATING EXPOSURE TO EMISSIONS | 6 |
| 2.1 | INTRODUCTION..... | 6 |
| 2.2 | POTENTIAL EXPOSURE PATHWAYS..... | 6 |
| 2.3 | EXPOSURE PATHWAYS CONSIDERED IN THE ASSESSMENT..... | 7 |
| 2.4 | EMISSIONS AND DISPERSION MODELLING INPUT DATA..... | 9 |
| 2.4.1 | <i>Compounds of Potential Concern (COPCs).....</i> | 9 |
| 2.4.2 | <i>Emission Parameters.....</i> | 10 |
| 2.4.3 | <i>Emission Concentrations for the COPCs.....</i> | 10 |
| 2.5 | DISPERSION MODELLING ASSUMPTIONS..... | 12 |
| 2.6 | DISPERSION MODELLING RESULTS..... | 13 |
| 3 | INPUT PARAMETERS FOR THE IRAP MODEL | 15 |
| 3.1 | INTRODUCTION..... | 15 |
| 3.2 | INPUT PARAMETERS FOR THE COPCs..... | 16 |
| 3.3 | SITE AND SITE SPECIFIC PARAMETERS..... | 17 |
| 3.4 | RECEPTOR INFORMATION..... | 18 |
| 4 | EXPOSURE ASSESSMENT | 20 |
| 4.1 | SELECTION OF RECEPTORS..... | 20 |
| 4.2 | ASSESSMENT OF INTAKE..... | 21 |
| 4.2.1 | <i>Ingestion Dose.....</i> | 21 |
| 4.2.2 | <i>Inhalation Dose.....</i> | 21 |
| 4.3 | EXPOSURE TO DIOXINS AND FURANS..... | 22 |
| 4.3.1 | <i>Comparison of Dioxin/Furan Exposure with WHO and UK COT Guidance.....</i> | 22 |
| 4.3.2 | <i>Infant Breast Milk Exposure to Dioxins and Furans.....</i> | 25 |
| 5 | SUMMARY AND CONCLUSIONS..... | 27 |
| 5.1 | SUMMARY..... | 27 |
| 5.2 | CONCLUSIONS..... | 27 |
| | | |
| ANNEX A | SITE PARAMETERS | |
| | | |
| ANNEX B | SCENARIO PARAMETERS | |


1 INTRODUCTION

1.1 BACKGROUND

Gair Consulting Ltd has been commissioned, on behalf of PyroCore Limited, by RPS to undertake an assessment to consider the effects on human exposure from emissions to air from two small waste incineration plants (SWIP) at an industrial site off Burcott Road in Avonmouth. The location of the installation is presented in *Figure 1.1*.

Incineration and co-incineration plant with a capacity to process less than 3 tonnes per hour of waste require a Schedule 13 SWIP permit to operate from the local authority, in this case Bristol City Council (BCC).

FIGURE 1.1 LOCATION OF THE PROPOSED INSTALLATION



1.2 DESCRIPTION OF THE INSTALLATION

1.2.1 Introduction

A small waste incineration (SWIP) permit for the operation of a Phoenix unit for the purpose of factory acceptance testing is held by Pyrocore. This HHRA has been prepared to support a variation to this permit to include a smaller demonstration unit on the same site.

1.2.2

Phoenix SWIP

PyroCore manufacture small waste incineration plants, more specifically, co-incineration plants. Prior to shipping to the Client's sites for operation, units may be installed and undergo factory acceptance tests and performance validation trials with the permitted wastes at a test site located at 203 Burcott Road, Avonmouth, Bristol, BS11 8AD.

Each SWIP to be tested will be of the same container sized footprint design (Phoenix model) and will be based on the processes of material pyrolysis and subsequent high temperature oxidation of the raw syngas. The Phoenix model has a capacity to process up to 500 - 600 kg/hr of waste, baselined on polyethylene terephthalate.

During factory acceptance testing the facility will initially be commissioned and tested using clean (virgin) biomass fuel before introducing other waste materials. Waste feed stocks will predominantly comprise plastic based wastes. The maximum annual throughput will be less than 200 tonnes of waste and 300 tonnes of biomass.

The Phoenix SWIP operates under the terms and conditions of a Schedule 13 permit which requires compliance with the ELVs for pollutants specified by the IED for small waste incineration plant. Effective pollutant abatement is achieved through the injection of a sodium bicarbonate reactant to achieve acid gas neutralisation and powdered activated carbon to abate potential dioxins, furans, volatile heavy metals and volatile organic compounds (VOCs). A ceramic filter facilitates removal of particulate bound heavy metals and other particulates and the reactants of the sodium bicarbonate, namely NaCl (sodium chloride), Na₂SO₄ (sodium sulphate). The Phoenix unit is already in place and is permitted to operate.

1.2.3

Demonstrator Unit

The demonstration (Demo) unit is also a SWIP and is based on pyrolysis and subsequent high temperature oxidation of raw syngas. PyroCore has operated its demonstration unit at Haybridge on a wide range of hazardous and non-hazardous waste materials and biomass from waste plastics to straw, grasses and woodchips from virgin sources.

The demonstration unit is to be relocated to the Avonmouth site where it will operate to conduct fully representative trials of customers own waste, or to confirm if the waste can be processed by PyroCore's pyrolysis technology. The demonstration unit has a capacity to process up to 250 kg per hour (kg/hr) of waste/fuel.

As per the Phoenix unit, the demonstration unit will comply with the emission limit values (ELVs) for pollutants specified by the Industrial Emissions

Directive (IED) for small waste incineration plant. The design of the air pollution control equipment is the same as the Phoenix unit.

1.3 PURPOSE OF THE ASSESSMENT

This Human Health Risk Assessment (HHRA) supplements the air quality assessment provided for the two SWIP. The HHRA only considers emissions to air as in this case human exposure to any harmful pollutants discharged directly to the aquatic environment and from solid waste disposal is considered to be negligible.

The area surrounding the installation is dominated by industrial land. Beyond the industrial area is the Severn Estuary to the west and agricultural areas to the east and northeast. The nearest populated areas are Shirehampton to the south (1.9 km) and Lawrence Weston to the southeast (2.3 km).

An air quality assessment of emissions from the installation has been provided by RPS¹. The air quality assessment provides a comparison of predicted concentrations of pollutants at off-site locations with background air quality and air quality standards and guidelines for the protection of human health. The air quality assessment assumes the theoretical position that the maximum permissible emission limit values (ELVs), stipulated for legal compliance for the two SWIP, are emitted during all times of operation. This position is considered unlikely to be a realistic operating scenario because, in reality, the emissions will be lower.

Given the above theoretical operating scenario, the emissions from the two SWIP would contain a number of substances that cannot be evaluated in terms of their effects on human health simply by reference to ambient air quality standards. Health effects could occur through exposure routes other than purely inhalation. As such, an assessment needs to be made of the overall human *exposure* to the substances by the local population and then the *risk* that this exposure causes.

This assessment has been undertaken to support the variation to the Environmental Permit for the installation and has been prepared in accordance with our understanding of the requirements of the Environment Agency for waste incineration and waste co-incineration plants. In the absence of local guidance, it is assumed that as the regulator, Bristol City Council's requirements will be similar. The Environment Agency requirements are for a human health risk assessment of dioxin/furan emissions from the installation based on the US EPA HHRAP methodology in the absence of UK or EU methods.

1 Air Quality Assessment - Proposed Small Waste Incineration Plant, 203 Burcott Road, Avonmouth, Bristol, July 2022 (Appendix B)

Human exposure to dioxins and furans has been compared against the Committee of Toxicity (COT) Tolerable Daily Intake (TDI) of 2 pg/kg per day. An assessment of exposure to dioxin-like polychlorinated biphenyls (PCBs) has also been included.

It should be noted that the former Her Majesty's Inspectorate of Pollution (HMIP) method does not have the capability to consider dioxin-like PCBs and the US EPA HHRAP method is limited in this respect. The HHRAP method does not contain physical properties or exposure parameters for individual dioxin-like PCBs but does provide information for two dioxin-like PCB mixtures (Aroclor 1016 and Aroclor 1254). Therefore, for these two substances typical emissions for dioxin-like PCBs have been included in the Industrial Risk Assessment Program (IRAP) model and these have been assumed to comprise entirely of Aroclor 1016 or Aroclor 1254 depending on which substance gives rise to the highest exposure.

1.4 SCOPE OF THE ASSESSMENT

An assessment needs to be made of the overall human *exposure* to the substances by the local population and then the *risk* that this exposure causes. The assessment presented here considers the potential impact of substances released by the installation on the health of the local population at the point of maximum exposure. These substances are those that are 'persistent' in the environment and have several pathways from the point of release to the human receptor. Essentially, they can be described as dioxins/furans and dioxin-like polychlorinated biphenyls (PCBs) and are present in extremely small quantities and are typically measured in mass units of nanograms ($\text{ng} = 10^{-9} \text{ g}$), picograms ($\text{pg} = 10^{-12} \text{ g}$) and femtograms ($\text{fg} = 10^{-15} \text{ g}$).

Unlike substances such as nitrogen dioxide, which have short term, acute effects on the respiratory system, dioxins/furans and dioxin-like PCBs have the potential if exposure is significant to cause effects through long term, cumulative exposure. A lifetime is the conventional period over which such effects are evaluated. A lifetime is taken to be 70 years.

The exposure scenarios used here represent highly unrealistic situations in which all exposure assumptions are chosen to represent a worst case and should be treated as an extreme view of the risks to health. While individual high-end exposure estimates may represent actual exposure possibilities (albeit at very low frequency), the possibility of all high end exposure assumptions accumulating in one individual is, for practical purposes, never realised. Therefore, intakes presented here should be regarded as an extreme upper theoretical representation of exposure that would be over and above that which would actually be experienced by the real population in the locality.

The risk assessment process is based on the application of the US EPA Human Health Risk Assessment Protocol (HHRAP)². This protocol has been assembled into a commercially available model, Industrial Risk Assessment Program (IRAP, Version 5.1.0) and marketed by Lakes Environmental of Ontario.

The approach seeks to quantify the *hazard* faced by the receptor, the *exposure* of the receptor to the substances identified as being a potential hazard and then to assess the *risk* of the exposure, as follows.

- *Quantification of the exposure*: an exposure evaluation determines the dose and intake of key indicator chemicals for an exposed person. The dose is defined as the amount of a substance contacting body boundaries (in the case of inhalation, the lungs) and intake is the amount of the substance absorbed into the body. The evaluation is based upon worst-case, conservative scenarios, with respect to the following:
 - location of the exposed individual and duration of exposure;
 - exposure rate;
 - emission rate from the source.
- *Risk characterisation*: following the above steps, the risk is characterised by examining the toxicity of the chemicals to which the individual has been exposed, and evaluating the significance of the calculated dose by a comparison of intakes with the tolerable daily intake (TDI) for dioxins/furans and dioxin-like PCBs.

2 US EPA Office of Solid Waste (September 2005) Human Health Risk Assessment Protocol for Hazardous Waste Combustion Facilities

2 METHODOLOGY FOR ESTIMATING EXPOSURE TO EMISSIONS

2.1 INTRODUCTION

An exposure assessment for the purposes of characterising the health impact of the proposed facility emissions requires the following steps:

- (1) Measurement or estimation of emissions from the source.
- (2) Modelling the fate and transport of the emitted substances through the atmosphere and through soil, water and biota following deposition onto land. Concentrations of the emitted chemicals in the environmental media are estimated at the point of exposure, which may be through inhalation or ingestion.
- (3) Calculation of the uptake of the emitted chemicals into humans coming into contact with the affected media and the subsequent distribution in the body.

With regard to Step (3), the exposure assessment considers the uptake of polychlorinated dibenzo-para-dioxins and polychlorinated dibenzofurans (PCDD/Fs, often abbreviated to 'dioxins/furans') and dioxin-like PCBs by various categories of human receptors.

2.2 POTENTIAL EXPOSURE PATHWAYS

There are two primary exposure 'routes' where humans may come into contact with chemicals that may be of concern:

- direct, via inhalation; or
- indirect, via ingestion of water, soil, vegetation and animals and animal products that become contaminated through the food chain.

There are four other potential exposure pathways of concern following the introduction of substances into the atmosphere:

- ingestion of drinking water;
- dermal (skin) contact with soil;
- incidental ingestion of soil; and
- dermal (skin) contact with water.


The possible exposure pathways included in the IRAP model are shown in *Figure 2.1*. Dermal contact with soil is an insignificant exposure pathway on the basis of the infrequent and sporadic nature of the events and the very low dermal absorption factors for this exposure route, coupled with the low plausible total dose that may be experienced (when considered over the lifetime of an individual). Health risk assessments of similar emissions (Pasternach (1989) *The Risk Assessment of Environmental and Human Health Hazards*, John Wiley, New York) have concluded that dermal absorption of soil is at least one order of magnitude less efficient than lung absorption.

Similar arguments are relevant with respect to the elimination of aquatic pathways from consideration; swimming, fishing and other recreational activities are also sporadic and unlikely to lead to significant exposures or uptake of any contamination into the human body via dermal contact with water.

Exposure via drinking water requires contamination of surface drinking water sources local to the point of consumption. The likelihood of contamination reaching a level of concern in the local water sources and ground water supplies is extremely low, particularly where there is no large-scale storage (e.g. reservoirs) or catchment areas for local water supplies. However, the US EPA's HHRAP does include the ingestion of drinking water from surface water sources as a potential exposure pathway where water bodies and water sheds have been defined within the exposure scenario. The ingestion of groundwater as a source of local drinking water is not considered by the HHRAP as it is considered to be an insignificant exposure pathway for emissions derived from combustion processes.

The ingestion of drinking water from surface water sources is only considered a potential exposure pathway where there is a local surface water body which provides local drinking water. However, it is our experience that drinking water from a reservoir located close to this type of facility makes a very small contribution to the total exposure. Therefore, exposure via drinking water is generally only considered where there is the potential for exposure via the ingestion of fish and the presence of edible fish farms (e.g. trout or salmon farms). There are no edible fish farms identified within 3 km of the proposed installation. The nearest coarse fishery (Green Park Fishing Lake) is located 4.5 km to the south-southeast. Furthermore, coarse fisheries are generally recreational fishing venues and coarse fish are not normally taken for consumption from these fisheries.

FIGURE 2.1 EXPOSURE PATHWAYS FOR RECEPTORS



On the basis of the assessment of the potential significance of the exposure pathways, the key exposure pathways which are relevant to the assessment and, hence, subject to examination in detail are as follows:

- inhalation;
- ingestion of food; and
- ingestion of soil.

Therefore, the exposures arising from ingestion are assessed with reference to the following:

- milk from home-reared cows;
- eggs from home-reared chickens;
- home-reared beef;
- home-reared pork;
- home-reared chicken;
- home-grown vegetable and fruit produce;
- breastmilk; and
- soil (incidental).

The inclusion of all food groups in the assessment conservatively assumes that both arable and pasture land are present in the vicinity of the predicted maximum annual average ground level concentration. This is, in reality, a highly unlikely scenario, but it has been included as a means of building a high degree of conservatism into the assessment and, hence, reducing the risk of exposures being underestimated. However, it should be noted that not all exposure scenarios will result in the ingestion of home-reared meat and animal products and these food products are only considered by the HHRAP for farmers and the families of farmers.

Similarly, the ingestion of fish is only considered where there is a local water body that is used for fishing and where the diet of the fisher (and family) may be regularly supplemented by fish caught from these local water sources. As discussed, there are no edible fish farms identified within 3 km of the proposed installation. Therefore, the ingestion of locally caught edible fish from an inland closed water source has not been considered as consumption rates are likely to be very small.

2.4 EMISSIONS AND DISPERSION MODELLING INPUT DATA

2.4.1 Compounds of Potential Concern (COPCs)

The substances which have been considered in the assessment are referred to as the Compounds of Potential Concern (COPCs) and include the seventeen PCDD/F congeners that are known to be toxic (refer *Section 2.4.3*). In addition,

the IRAP model includes two dioxin-like PCBs (Aroclor 1016 and Aroclor 1254). These comprise a mixture of congeners with one to four chlorine atoms for Aroclor 1016 with a chlorine content of 41% by mass (average of three chlorine atoms). Similarly, Aroclor 1254 has between four and seven chlorine atoms and a chlorine content of 54% by mass (average of five chlorine atoms).

2.4.2 Emission Parameters

Emissions from the two SWIP will be via separate stacks. Emission parameters assumed for the assessment are consistent with those used for the air quality and for the Demo SWIP are as follows:

- stack height of 11 m (metres) above ground level;
- flue diameter of 0.4 m;
- emission temperature of 200°C (degrees celcius) or 473 K (kelvin);
- emission velocity of 10.0 m s⁻¹ (metres per second); and
- normalised flow rate of 0.44 Nm³ s⁻¹ (normal cubic metres per second at 273 K, dry and 11% O₂).

For the Phoenix SWIP the emission parameters are as follows:

- stack height of 10 m above ground level;
- flue diameter of 0.5 m;
- emission temperature of 250°C or 523 K;
- emission velocity of 19.9 m s⁻¹; and
- normalised flow rate of 1.96 Nm³ s⁻¹.

2.4.3 Emission Concentrations for the COPCs

The general term dioxins denotes a family of compounds, with each compound composed of two benzene rings interconnected with two oxygen atoms. There are 75 individual dioxins, with each distinguished by the position of chlorine or other halogen atoms positioned on the benzene rings. Furans are similar in structure to dioxins, but have a carbon bond instead of one of the two oxygen atoms connecting the two benzene rings. There are 135 individual furan compounds. Each individual furan or dioxin compound is referred to as a congener and each has a different toxicity and physical properties with regard to its atmospheric behaviour. It is important, therefore, that the exposure methodology determines the fate and transport of PCDD/Fs on a congener specific basis. It does this by accounting for the varying volatility of the congeners and their different toxicities. Consequently, information regarding the PCDD/F annual mean ground level concentrations on a congener specific basis is required.

For the purposes of the exposure assessment, the congener profile for the two proposed SWIP is presented in *Table 2.1*, which is a standard profile for municipal waste incinerators derived by Her Majesty's Inspectorate of Pollution (HMIP), one of the predecessors of the Environment Agency. The international toxic equivalency factors are given and used to derive the toxic equivalent emission (I-TEQ). It is assumed that PCDD/F emissions are 0.1 ng I-TEQ Nm⁻³ (reference conditions 273K, dry and 11% O₂). This assumption is justified because it is derived from the IED emission limit value which will be controlled and enforced by the conditions of the Schedule 13 permit.

TABLE 2.1 PCDD/F CONGENER PROFILE FOR THE PROPOSED SWIP

| Congener | Annual Mean Emission Concentration (ng Nm ⁻³) (a) | I-TEF toxic equivalent factors) | Annual Mean Emission Concentration (b) (ng I-TEQ Nm ⁻³) |
|--|---|---------------------------------|---|
| 2,3,7,8-TCDD | 0.0031 | 1.0 | 0.0031 |
| 1,2,3,7,8-PeCDD | 0.025 | 0.5 | 0.012 |
| 1,2,3,4,7,8-HxCDD | 0.029 | 0.1 | 0.0029 |
| 1,2,3,7,8,9-HxCDD | 0.021 | 0.1 | 0.0021 |
| 1,2,3,6,7,8-HxCDD | 0.026 | 0.1 | 0.0026 |
| 1,2,3,4,6,7,8-HpCDD | 0.17 | 0.01 | 0.0017 |
| OCDD | 0.40 | 0.001 | 0.00040 |
| 2,3,7,8-TCDF | 0.028 | 0.1 | 0.0028 |
| 2,3,4,7,8-PeCDF | 0.054 | 0.5 | 0.027 |
| 1,2,3,7,8-PeCDF | 0.028 | 0.05 | 0.0014 |
| 1,2,3,4,7,8-HxCDF | 0.22 | 0.1 | 0.022 |
| 1,2,3,7,8,9-HxCDF | 0.0040 | 0.1 | 0.00040 |
| 1,2,3,6,7,8-HxCDF | 0.081 | 0.1 | 0.0081 |
| 2,3,4,6,7,8-HxCDF | 0.087 | 0.1 | 0.0087 |
| 1,2,3,4,6,7,8-HpCDF | 0.44 | 0.01 | 0.0044 |
| 1,2,3,4,7,8,9-HpCDF | 0.040 | 0.01 | 0.00040 |
| OCDF | 0.40 | 0.001 | 0.00040 |
| Total (ng I-TEQ m⁻³) | | | 0.1 |
| (a) Congener profile from Table 7.2a DOE (1996) Risk Assessment of Dioxin Releases from Municipal Waste Incineration Processes Contract No. HMIP/CPR2/41/1/181 | | | |
| (b) Reference conditions of 273K, 1 atmosphere, dry and 11% O ₂ | | | |

In accordance with the Air Quality Assessment ¹, the emission concentration for PCBs is assumed to be 8 x 10⁻⁸ mg m⁻³. As a worst-case it is assumed to comprise entirely of dioxin-like PCBs.

The emission rates for each substance as input to the IRAP model are provided in *Table 2.2*. It is assumed that the Demo SWIP operates continuously. However, the Phoenix SWIP is only expected to run continuously for two

months per year and for the remaining ten months only for one week per month. Therefore, it will only operate for 18 weeks a year. Therefore, emissions for the Phoenix SWIP have been adjusted by a factor of 18/52 to account for the reduced operational hours.

TABLE 2.2 PCDD/F EMISSION RATES USED IN THE IRAP MODEL

| Congener | Emission Concentration (mg Nm ⁻³) | Demo SWIP Emission Rate (g s ⁻¹) | Phoenix SWIP Emission Rate (g s ⁻¹) |
|---------------------|---|--|---|
| 2,3,7,8-TCDD | 0.0031 x 10 ⁻⁶ | 1.4 x 10 ⁻¹² | 2.1 x 10 ⁻¹² |
| 1,2,3,7,8-PeCDD | 0.025 x 10 ⁻⁶ | 1.1 x 10 ⁻¹¹ | 1.7 x 10 ⁻¹¹ |
| 1,2,3,4,7,8-HxCDD | 0.029 x 10 ⁻⁶ | 1.3 x 10 ⁻¹¹ | 2.0 x 10 ⁻¹¹ |
| 1,2,3,7,8,9-HxCDD | 0.021 x 10 ⁻⁶ | 9.2 x 10 ⁻¹² | 1.4 x 10 ⁻¹¹ |
| 1,2,3,6,7,8-HxCDD | 0.026 x 10 ⁻⁶ | 1.1 x 10 ⁻¹¹ | 1.8 x 10 ⁻¹¹ |
| 1,2,3,4,6,7,8-HpCDD | 0.17 x 10 ⁻⁶ | 7.5 x 10 ⁻¹¹ | 1.2 x 10 ⁻¹⁰ |
| OCDD | 0.40 x 10 ⁻⁶ | 1.8 x 10 ⁻¹⁰ | 2.7 x 10 ⁻¹⁰ |
| 2,3,7,8-TCDF | 0.028 x 10 ⁻⁶ | 1.2 x 10 ⁻¹¹ | 1.9 x 10 ⁻¹¹ |
| 2,3,4,7,8-PeCDF | 0.054 x 10 ⁻⁶ | 2.4 x 10 ⁻¹¹ | 3.6 x 10 ⁻¹¹ |
| 1,2,3,7,8-PeCDF | 0.028 x 10 ⁻⁶ | 1.2 x 10 ⁻¹¹ | 1.9 x 10 ⁻¹¹ |
| 1,2,3,4,7,8-HxCDF | 0.22 x 10 ⁻⁶ | 9.6 x 10 ⁻¹¹ | 1.5 x 10 ⁻¹⁰ |
| 1,2,3,7,8,9-HxCDF | 0.0040 x 10 ⁻⁶ | 1.8 x 10 ⁻¹² | 2.7 x 10 ⁻¹² |
| 1,2,3,6,7,8-HxCDF | 0.081 x 10 ⁻⁶ | 3.6 x 10 ⁻¹¹ | 5.5 x 10 ⁻¹¹ |
| 2,3,4,6,7,8-HxCDF | 0.087 x 10 ⁻⁶ | 3.8 x 10 ⁻¹¹ | 5.9 x 10 ⁻¹¹ |
| 1,2,3,4,6,7,8-HpCDF | 0.44 x 10 ⁻⁶ | 1.9 x 10 ⁻¹⁰ | 3.0 x 10 ⁻¹⁰ |
| 1,2,3,4,7,8,9-HpCDF | 0.040 x 10 ⁻⁶ | 1.8 x 10 ⁻¹¹ | 2.7 x 10 ⁻¹¹ |
| OCDF | 0.40 x 10 ⁻⁶ | 1.8 x 10 ⁻¹⁰ | 2.7 x 10 ⁻¹⁰ |
| Aroclor 1016/1254 | 0.080 x 10 ⁻⁶ | 3.5 x 10 ⁻¹¹ | 5.4 x 10 ⁻¹¹ |

2.5 DISPERSION MODELLING ASSUMPTIONS

The air quality assessment supporting the permit application has relied upon the use of ADMS to estimate ground level concentrations of pollutants. The HHRA model has been designed to accept output files from the US EPA ISC or AERMOD dispersion models, reflecting its North American origins and its need to follow the US EPA risk assessment protocol. The use of ADMS is consistent with the air quality assessment undertaken for the two SWIP and the emissions data and model set up are identical to that carried out for the air quality assessment ¹.

Therefore, to maintain consistency with the air quality assessment, it has been possible to use output from the ADMS model with IRAP using the following procedure:

- generation of ISC input files and output files for the study area;
- generation of ADMS output data using the approach outlined in the US EPA risk assessment protocol; and
- inserting the ADMS results into the ISC output files.

For the modelling, all emission properties, building heights, and other relevant factors were retained from the air quality assessment provided for the installation. As the health risk assessment requires information on the deposition of substances to surfaces as well as airborne concentrations of substances, the ADMS dispersion model has also been used to predict the following:

- the airborne concentration of vapour, particle and particle bound substances emitted;
- the wet deposition rate of particle and particle bound substances; and
- the dry deposition rate of vapour, particle and particle bound substances.

For dry deposition of particles and particle bound contaminants a fixed deposition velocity of 0.01 m s^{-1} has been used. The two SWIP will be equipped with ceramic filters for particle removal and the emitted particles are likely to be less than $1 - 2 \mu\text{m}$ in diameter. For particles of this size, deposition velocities are likely to be of the order of 0.001 to 0.01 m s^{-1} . Therefore, as a worst-case, for the ADMS modelling a value of 0.01 m s^{-1} has been adopted. A gas dry deposition velocity of 0.005 m s^{-1} is used for the gas phase. For wet deposition, the following washout coefficients are used:

- Gas phase - washout coefficient A at 0.00016 and washout coefficient B of 0.64;
- Particle phase - washout coefficient A at 0.00028 and washout coefficient B of 0.64; and
- Particle bound phase - washout coefficient A at 0.00010 and washout coefficient B of 0.64.

2.6 DISPERSION MODELLING RESULTS

A summary of the key results from the ADMS dispersion model is presented in *Table 2.3*. These have been predicted using the 2017 Filton meteorological data set. This year was selected, as out of the five years considered (2013 to 2017), it was the year that provided highest predicted annual mean concentrations and deposition rates.

TABLE 2.3

MAXIMUM ANNUAL AVERAGE PARTICLE PHASE CONCENTRATIONS AND PARTICLE PHASE DEPOSITION RATES ESTIMATED BY ADMS

| Pollutant | Max Annual Average Concentration (a) (fg m ⁻³) | | Max Annual Average Deposition Rate (b) (ng m ⁻² year ⁻¹) | |
|---|---|--------------|--|--------------|
| | Demo SWIP | Phoenix SWIP | Demo SWIP | Phoenix SWIP |
| 2,3,7,8-TCDD | 0.049 | 0.10 | 0.016 | 0.032 |
| 1,2,3,7,8-PeCDD | 0.39 | 0.82 | 0.13 | 0.25 |
| 1,2,3,4,7,8-HxCDD | 0.46 | 0.97 | 0.15 | 0.30 |
| 1,2,3,7,8,9-HxCDD | 0.33 | 0.70 | 0.11 | 0.21 |
| 1,2,3,6,7,8-HxCDD | 0.41 | 0.87 | 0.14 | 0.27 |
| 1,2,3,4,6,7,8-HpCDD | 2.7 | 5.7 | 0.88 | 1.7 |
| OCDD | 6.3 | 13.4 | 2.1 | 4.1 |
| 2,3,7,8-TCDF | 0.44 | 0.94 | 0.15 | 0.29 |
| 2,3,4,7,8-PeCDF | 0.85 | 1.8 | 0.28 | 0.55 |
| 1,2,3,7,8-PeCDF | 0.44 | 0.94 | 0.15 | 0.29 |
| 1,2,3,4,7,8-HxCDF | 3.4 | 7.3 | 1.1 | 2.2 |
| 1,2,3,7,8,9-HxCDF | 0.063 | 0.13 | 0.021 | 0.041 |
| 1,2,3,6,7,8-HxCDF | 1.3 | 2.7 | 0.42 | 0.83 |
| 2,3,4,6,7,8-HxCDF | 1.4 | 2.9 | 0.45 | 0.89 |
| 1,2,3,4,6,7,8-HpCDF | 7.0 | 14.8 | 2.3 | 4.5 |
| 1,2,3,4,7,8,9-HpCDF | 0.63 | 1.3 | 0.21 | 0.41 |
| OCDF | 6.3 | 13.4 | 2.1 | 4.1 |
| Aroclor 1016/1254 | 1.3 | 2.7 | 0.42 | 0.82 |
| (a) Where 1 fg m ⁻³ is equal to 1 x 10 ⁻¹⁵ g m ⁻³ | | | | |
| (b) Where 1 ng m ⁻² year ⁻¹ is equal to 1 x 10 ⁻⁹ g m ⁻² year ⁻¹ | | | | |

3.1 INTRODUCTION

Exposure of an individual to a chemical may occur either by inhalation or ingestion (including food, water and soil). Of interest is the total dose of the chemical received by the individual through the combination of possible routes, and the IRAP model has been developed to estimate the dose received by the human body, often referred to as the external dose.

Exposure to COPCs is a function of the estimated concentration of the substance in the environmental media with which individuals may come into contact (i.e. exposure point concentrations) and the duration of contact. The concentration at the point of contact is itself a function of the transfer through air, soil, water, plants and animals that form part of the overall pathway. Exposure equations have been developed which combine exposure factors (e.g. exposure duration, frequency and medium intake rate) and exposure point concentrations. The dose equations therefore facilitate estimation of the received dose and account for the properties of the route of exposure, i.e. ingestion and inhalation.

For those substances that bio-accumulate, i.e. become more concentrated higher up the food chain, especially in body fats, the exposure to contaminated meat products and milk is of particular significance.

The IRAP model user has the facility to adjust some of the key exposure factors. An example is the diet of the receptor and the proportion of which is local produce, which may be contaminated. Obviously, if a nearby resident eats no food grown locally, then that person's diet cannot be contaminated by the emissions from the source, in this case the proposed facility. It is conventional to investigate two types of receptor, a farmer and a resident. It is assumed that a farmer eats proportionately more locally grown food than a resident. Where the potential exists for the consumption of locally caught fish a fisher receptor may also be considered.

The receptor types can also be divided into adults and children. Children are important receptors because they tend to ingest soil and dusts directly and have lower body weights, so that the effect of the same dose is greater in the child than in the adult.

The IRAP model is designed to accept output files of airborne concentrations and deposition rates. From these, it proceeds to calculate the concentrations of the pollutants of concern in the environmental media, foodstuffs and the human receptor. The dose experienced by the human receptor can be compared to the tolerable daily intake (TDI) provided by the Committee on Toxicity for dioxins and dioxin like PCBs of $2 \text{ pg kg}^{-1} \text{ d}^{-1}$.

The model requires a wide range of input parameters to be defined, these include:

- physical and chemical properties of the COPCs;
- site information, including site specific data; and
- receptor information – for each receptor type (e.g. adult or child, resident or farmer or fisher).

The HHRAP default values, which are incorporated into the IRAP model, have been used for the majority of these input values. These data are provided in the following sections.

3.2 INPUT PARAMETERS FOR THE COPCS

The IRAP model contains a database of physical and chemical parameters for each of the 206 COPCs. This database is based on default values provided by the HHRAP and all default values have been used for this assessment.

These parameters are used to determine how each of the COPCs behave in the environment and their presence and accumulation in various food products (meat, fish, animal products, vegetation, soil and water). For 2,3,7,8-TCDD (the most toxic of the PCDD/Fs), the default parameters are provided in *Table 3.1*.

TABLE 3.1 IRAP INPUT PARAMETERS FOR 2, 3, 7, 8-TCDD

| Parameter Description | Symbol | Units | 2,3,7,8-TCDD |
|--|---------|--------------------------------------|--------------------------|
| Chemical abstract service number | CAS No. | - | 1746-01-6 |
| Molecular weight | MW | g mole ⁻¹ | 322.0 |
| Melting point of chemical | T_m | K | 578.7 |
| Vapour pressure | V_p | atm | 1.97 x 10 ⁻¹² |
| Aqueous solubility | S | mg L ⁻¹ | 1.93 x 10 ⁻⁵ |
| Henry's Law constant | H | atm-m ³ mol ⁻¹ | 3.29 x 10 ⁻⁵ |
| Diffusivity of COPC in air | D_a | cm ² s ⁻¹ | 0.104 |
| Diffusivity of COPC in water | D_w | cm ² s ⁻¹ | 5.6 x 10 ⁻⁶ |
| Octanol-water partition coefficient | K_ow | - | 6,309,573 |
| Organic carbon-water partition coefficient | K_oc | mL g ⁻¹ | 3,890,451 |
| Soil-water partition coefficient | Kd_s | mL g ⁻¹ | 38,904 |
| Suspended sediments/surface water partition coefficient | Kd_sw | L kg ⁻¹ | 291,784 |
| Bed sediment/sediment pore water partition coefficient | Kd_bs | mL g ⁻¹ | 155,618 |
| COPC loss constant due to biotic and abiotic degradation | K_sg | a ⁻¹ | 0.03 |
| Fraction of COPC air concentration in vapour phase | f_v | | 0.664 |
| Root concentration factor | RCF | mL g ⁻¹ | 39,999 |

TABLE 3.1 IRAP INPUT PARAMETERS FOR 2, 3, 7, 8-TCDD

| Parameter Description | Symbol | Units | 2,3,7,8-TCDD |
|---|--------------|----------------------|--------------|
| Plant-soil bioconcentration factor for below ground produce | br_root_veg | - | 1.03 |
| Plant-soil bioconcentration factor for leafy vegetables | br_leafy_veg | - | 0.00455 |
| Plant-soil bioconcentration factor for forage | br_forage | - | 0.00455 |
| COPC air-to-plant biotransfer factor for leafy vegetables | bv_leafy_veg | - | 65,500 |
| COPC air-to-plant biotransfer factor for forage | bv_forage | - | 65,500 |
| COPC biotransfer factor for milk | ba_milk | day kg ⁻¹ | 0.0055 |
| COPC biotransfer factor for beef | ba_beef | day kg ⁻¹ | 0.026 |
| COPC biotransfer factor for pork | ba_pork | day kg ⁻¹ | 0.032 |
| Bioconcentration factor for COPC in eggs | Bcf_egg | - | 0.060 |
| Bioconcentration factor for COPC in chicken | Bcf_chicken | - | 3.32 |
| Fish bioconcentration factor | BCF_fish | L kg ⁻¹ | 34,400 |
| Fish bioaccumulation factor | BAF_fish | L kg ⁻¹ | 0 |
| Biota-sediment accumulation factor | BSAF_fish | - | 0.09 |
| Plant-soil bioconcentration factor for grain | br_grain | - | 0.00455 |
| Plant-soil bioconcentration factor for eggs | br_egg | - | 0.011 |
| COPC biotransfer factor for chicken | ba_chicken | day kg ⁻¹ | 0.019 |

3.3 SITE AND SITE SPECIFIC PARAMETERS

The IRAP health risk assessment model requires information relating to the location and its surroundings. The parameters required include the following.

- The fraction of animal feed (grain, silage and forage) grown on contaminated soils and quantity of animal feed and soil consumed by the various animal species considered.
- The interception fraction for above ground vegetation, forage and silage and length of vegetation exposure to deposition. The yield/standing crop biomass is also required.
- Input data for assessing the risks associated with exposure to breast milk, including:
 - body weight of infant;
 - exposure duration;
 - proportion of ingested COPC stored in fat;
 - proportion of mother's weight that is fat;

- fraction of fat in breast milk;
- fraction of ingested contaminant that is absorbed; and
- half-life of dioxins in adults and ingestion rate of breast milk.
- Other physical parameters (e.g. soil dry bulk density, density of air, soil mixing zone depth).

For all of these parameters the IRAP/EPA HHRAP default values have been used and these are presented in *Annex A*. Other site specific parameters are also required which are not provided by the IRAP model. These parameters were specified for the proposed facility as follows:

- Annual average evapotranspiration rate of 53.3 cm a⁻¹ (assumed to be 70% of total precipitation);
- Annual average precipitation of 76.2 cm a⁻¹ (based on the average for the five year data set for the 2013 to 2017 meteorological data);
- Annual average irrigation of 0 cm a⁻¹ since manual irrigation of crops in the UK is not generally required due to natural irrigation;
- Annual average runoff of 7.6 cm a⁻¹ (assumed to be 10% of total precipitation);
- An annual average wind velocity of 4.2 m s⁻¹ (average for the five years); and
- A time period over which deposition occurs of 30 years (the HHRAP default value).

3.4 RECEPTOR INFORMATION

Within the IRAP model there are three receptor types; Resident, Farmer and Fisher. Information relating to each receptor type (adult and/or child) is required by the model where these receptor types are used. The information required includes the following:

- Food (meat, dairy products, fish and vegetables), water and soil consumption rates for each receptor type. However, only Fishers are assumed to consume fish and only Farmers are assumed to consume locally reared animals and animal products.
- Fraction of contaminated food, water and soil which is consumed by each receptor type.
- Input data for the inhalation exposure including: inhalation exposure duration, inhalation exposure frequency, inhalation exposure time; and inhalation rate.
- Input data for the ingestion exposure including: exposure duration, exposure frequency, exposure time; and body weight of receptor.

For the purposes of this assessment the default IRAP/HHRAP parameters have been used mainly to define the characteristics of the receptors. The default input data are presented in *Annex B*. The only variation to this is the assumed body weight of a child receptor. The IRAP/HHRAP default value is 15 kg whereas in the UK a value of 20 kg is typically used. Therefore, a value of 20 kg has been used.

4.1 SELECTION OF RECEPTORS

In addition to defining specific locations for assessment, IRAP can be used to determine the location of the maximum impact over an area based on the results of the dispersion model. For each defined land-use area, IRAP selects the locations which represent the maximum predicted concentrations or deposition rates for the area selected. The locations of these various maxima are often co-located resulting in the selection of one to nine receptor locations per defined area. This approach is adopted by IRAP since the maximum receptor impact may occur at any one of the maximum concentration or deposition locations identified.

The nearest residential areas comprise residential properties in Shirehampton to the south and Lawrence Weston to the southeast. Therefore, two areas where residential exposure may occur have been defined.

The nearest agricultural land to the site is to the southeast but is likely to be of poor quality. Two areas where the potential for farming exists have been defined. These include areas to the northeast (NE) and southeast (SE).


For each type of receptor up to nine locations are selected based on the maximum predicted airborne concentration, maximum predicted wet deposition rate and maximum dry deposition rate for the gas phase, particle phase and particle bound phase. For the assessment, three Farmer receptors and three Residential receptors have been assessed. It is considered that the likelihood of locally caught fish being consumed is low and fisher receptors have not been included in the assessment. For all of the receptor types, adult and child receptors have been considered. The locations of the Resident and Farmer receptors are described in *Table 4.1* and presented in *Figure 4.1*.

TABLE 4.1 DESCRIPTION OF RESIDENT AND FARMER RECEPTORS

| Ref. | Name | Type | Easting | Northing |
|------|----------------------------|----------|---------|----------|
| FNE | Farmer NE | Farmer | 353539 | 180582 |
| FSE1 | Farmer SE 1 | Farmer | 353889 | 180182 |
| FSE2 | Farmer SE 2 | Farmer | 352689 | 179682 |
| RLW1 | Resident Lawrence Weston 1 | Resident | 354489 | 178982 |
| RLW2 | Resident Lawrence Weston 2 | Resident | 354539 | 178982 |
| RS | Resident Shirehampton | Resident | 351689 | 178432 |

At other locations not specifically considered in the assessment, the predicted hazards and risks will be lower than predicted for the discrete receptors considered.

FIGURE 4.1 LOCATION OF THE RESIDENT AND FARMER RECEPTORS



4.2 ASSESSMENT OF INTAKE

4.2.1 Ingestion Dose

The ingestion intake is calculated as the Average Daily Dose (ADD) from all ingestion exposure routes (e.g. soil, above ground vegetables, meat and dairy products) where for example:

$$ADD_{Ing, TCDD} = \frac{I_{Ing, TCDD} \bullet ED \bullet EF}{AT \bullet 365}$$

Where: $ADD_{Ing, TCDD}$ = total ingestion dose for TCDD; ED is the exposure duration (dependent on the receptor type); EF is the exposure frequency (350 days per year); and AT is the averaging time, and for determining the TDI, is assumed to be equal to the ED. The total dose is the sum of the dose for each of the individual congeners.

4.2.2 Inhalation Dose

For inhalation, the ADD from inhalation exposure is calculated as follows:

$$ADD_{Inh, TCDD} = \frac{C_a \bullet IR \bullet ED \bullet EF}{AT \bullet 365}$$

Where: $ADD_{Inh, TCDD}$ is the total inhalation dose for TCDD, C_a is the concentration of TCDD in air and IR is the daily inhalation rate. The total dose is the sum of the dose for each of the individual congeners.

4.3 EXPOSURE TO DIOXINS AND FURANS

4.3.1 Comparison of Dioxin/Furan Exposure with WHO and UK COT Guidance

Facility Contribution to Intake

The World Health Organization (WHO) recommends a tolerable daily intake for dioxins/furans of 1 to 4 pg I-TEQ kg-BW⁻¹ d⁻¹ (picogrammes as the International Toxic Equivalent per kilogram bodyweight per day)³. The TDI represents the tolerable daily intake for lifetime exposure and short-term excursions above the TDI would have no consequence provided that the average intake over long periods is not exceeded. The average (lifetime) daily intake of dioxins/furans for the receptors considered is presented in *Table 4.2* for the individual emissions from each SWIP. These are also compared to the Committee on Toxicity (COT) TDI for dioxins and dioxin-like PCBs of 2 pg I-TEQ kg-BW⁻¹ d⁻¹.

TABLE 4.2 COMPARISON OF AVERAGE DAILY INTAKES WITH THE UK COT AND WHO'S TDI FOR DIOXINS/FURANS FOR THE DEMO SWIP AND PHOENIX SWIP (pg I-TEQ kg-BW⁻¹ d⁻¹)

| Receptor Name | Demo SWIP | | Phoenix SWIP | |
|----------------------------|---|----------|--------------|----------|
| | Adult | Child | Adult | Child |
| Farmer NE | 0.0032 | 0.0046 | 0.0038 | 0.0055 |
| Farmer SE 1 | 0.0016 | 0.0023 | 0.0019 | 0.0028 |
| Farmer SE 2 | 0.0012 | 0.0017 | 0.0014 | 0.0021 |
| Resident Lawrence Weston 1 | 0.0000086 | 0.000024 | 0.0000086 | 0.000024 |
| Resident Lawrence Weston 2 | 0.0000085 | 0.000024 | 0.0000085 | 0.000024 |
| Resident Shirehampton | 0.0000087 | 0.000024 | 0.000010 | 0.000029 |
| WHO TDI | 1 to 4 pg I-TEQ kg-BW ⁻¹ d ⁻¹ | | | |
| Committee on Toxicity | 2 pg I-TEQ kg-BW ⁻¹ d ⁻¹ | | | |

The contribution of the Phoenix SWIP to the average daily intake is up to 120% higher than for the Demo SWIP. Although the Demo SWIP is a smaller emission source it is assumed to operated continuously whereas the Phoenix SWIP is only assumed to operate for 18 weeks per year.

The average (lifetime) daily intake of dioxins/furans for the combined Demo and Phoenix SWIP is presented in *Table 4.3*.

3 Assessment of the Health Risk of Dioxins: Re-evaluation of the Tolerable Daily Intake (TD), WHO Consultation, May 25-29 1998, Geneva, Switzerland

TABLE 4.3

COMPARISON OF AVERAGE DAILY INTAKES WITH THE UK COT AND WHO'S TDI FOR DIOXINS/FURANS FOR THE COMBINED EMISSIONS FROM THE DEMO SWIP AND PHOENIX SWIP ($\mu\text{g I-TEQ kg-BW}^{-1} \text{ d}^{-1}$)

| Receptor Name | Demo and Phoenix SWIP Combined | |
|----------------------------|--|----------|
| | Adult | Child |
| Farmer NE | 0.0070 | 0.010 |
| Farmer SE 1 | 0.0035 | 0.0051 |
| Farmer SE 2 | 0.0026 | 0.0038 |
| Resident Lawrence Weston 1 | 0.000017 | 0.000048 |
| Resident Lawrence Weston 2 | 0.000017 | 0.000048 |
| Resident Shirehampton | 0.000019 | 0.000053 |
| WHO TDI | 1 to 4 $\mu\text{g I-TEQ kg-BW}^{-1} \text{ d}^{-1}$ | |
| Committee on Toxicity | 2 $\mu\text{g I-TEQ kg-BW}^{-1} \text{ d}^{-1}$ | |

The maximum contribution of the two SWIP to the COT TDI is 0.5% for the Farmer Northeast child receptor and 0.3% for the Farmer Northeast adult receptor. This assumes as a worst-case that these receptors produce their own home reared and home-grown food at the location of maximum impact for the area and represents an extreme worst-case. Furthermore, this assumes that both arable land and pastureland are available at this location. Therefore, it is considered that the predicted impacts for this receptor and for other farmer receptors represent an extreme worst-case.

For the residential receptors, the maximum contribution of the two SWIP to the COT TDI is <0.1% for Resident Shirehampton. Therefore, the contribution of the facility to the intake of dioxins/furans and dioxin-like PCBs is negligible.

Total Intake

The contribution of the facility to total intake is provided as follows:

- predicted incremental intake due to emissions from the facility;
- average daily background intake (i.e. that arising from other sources), referred to as the mean daily intake (MDI);
- the total intake (i.e. the sum of the predicted incremental intake and the MDI);
- a comparison of the total intake with the TDI for dioxin/furans.

For the key receptors (i.e. those which represent the predicted highest exposure for the receptor types considered) the results are presented in *Table 4.4*. Results are presented for Farmer Northeast and Resident Shirehampton where highest farmer and resident exposures are predicted. Results are presented for both adult and child receptors.

The MDI is derived from data provided by the Environment Agency ⁴ and a value of 49 pg WHO-TEQ d⁻¹. The MDI for an adult receptor and child receptor is calculated as follows:

- for an adult receptor a MDI of 0.7 pg I-TEQ kg⁻¹ d⁻¹ ⁵ is derived by dividing the Environment Agency MDI by a bodyweight of 70 kg;
- for a child receptor a MDI of 1.8 pg I-TEQ kg⁻¹ d⁻¹ is derived by dividing the Environment Agency MDI by a bodyweight of 20 kg and applying an adult to child correction factor of 0.74.

TABLE 4.4 COMPARISON OF TOTAL INTAKE WITH THE COT TDI FOR THE COMBINED EMISSIONS

| Receptor | Total Intake from the Facility (pg I-TEQ kg ⁻¹ d ⁻¹) | Total Intake Facility + MDI (pg I-TEQ kg ⁻¹ d ⁻¹) | Facility as %age of TDI | Total Intake as %age of TDI |
|-----------------------------|---|--|-------------------------|-----------------------------|
| Farmer Northeast Adult | 0.0070 | 0.71 | 0.3% | 35.3% |
| Farmer Northeast Child | 0.010 | 1.81 | 0.5% | 90.5% |
| Resident Shirehampton Adult | 0.000019 | 0.70 | <0.1% | 35.0% |
| Resident Shirehampton Child | 0.000053 | 1.80 | <0.1% | 90.0% |
| COT TDI | 2 | 2 | - | - |

For inhalation and oral intake of PCDD/Fs for adults, total intake is well below the TDI. Background exposure represents approximately 35% of total exposure. At worst, the two SWIP contribute 0.3% to the TDI for adults.

For inhalation and oral intake of PCDD/Fs for children, the background intake is relatively high at 90% of the TDI. At worst, the additional contribution from the two SWIP for a child is 0.010 pg TEQ kg⁻¹ d⁻¹ (0.5% of the COT TDI). Combined with the background exposure for a 20 kg child (1.8 pg TEQ kg⁻¹ d⁻¹) the total intake would be below the TDI (90.5%). Furthermore, it should be noted that the TDI for PCCD/Fs is set for the purposes of assessing lifetime exposure and these elevated background exposures for children are not representative of long-term exposure. Therefore, taking into account the extreme worst-case assumptions adopted for farmer receptors, it is concluded that the contribution of the two SWIP to total intake would be not significant.

4 Soil Guideline Values for dioxins, furans and dioxin-like PCBs in soil, Environment Agency, Science Report SC050021/Dioxins SGV, September 2009

5 No correction is provided between the WHO-TEF and the I-TEF but a sensitivity analysis indicates that correcting between the two systems would have negligible impact on the results

4.3.2

Infant Breast Milk Exposure to Dioxins and Furans

Another exposure pathway of interest is infant exposure to dioxins and furans via the ingestion of their mother's breast milk. This is because the potential for contamination of breast milk is particularly high for dioxin-like compounds such as these, as they are extremely lipophilic (fat soluble) and hence likely to accumulate in breast milk. Further, the infant body weight is smaller and it could be argued that the effect is proportionately greater than in an adult. However, this would only be the case if the dioxin-like emissions were significant.

This exposure is measured by the Average Daily Dose (ADD) on the basis of an averaging time of 1 year. In the US, a threshold value of 50 pg kg⁻¹ d⁻¹ of 2,3,7,8-TCDD TEQ is cited as being potentially harmful. The IRAP model calculates the ADD that would result from an adult receptor breast feeding an infant. It should be noted that the ADD from breast feeding calculated by IRAP does not consider dioxin-like PCBs. However, the dioxin-like PCB emission is a small fraction of the total emission and the inclusion of dioxin-like PCBs would not result in a significant increase in the ADD from breast feeding.

For the combined SWIP emissions, a summary of the ADD for each of the infants of adult receptors considered for the assessment is presented in *Table 4.5*.

TABLE 4.5 ASSESSMENT OF THE AVERAGE DAILY DOSE FOR A BREAST-FED INFANT OF AN ADULT RECEPTOR – COMBINED EMISSIONS

| Receptor Name | Average Daily Dose from Breast Feeding (pg kg ⁻¹ d ⁻¹ of 2,3,7,8-TCDD) |
|----------------------------|---|
| Farmer NE | 0.082 |
| Farmer SE 1 | 0.041 |
| Farmer SE 2 | 0.030 |
| Resident Lawrence Weston 1 | 0.00016 |
| Resident Lawrence Weston 2 | 0.00016 |
| Resident Shirehampton | 0.00018 |
| <i>US EPA Criterion</i> | <i>50</i> |
| <i>WHO criterion</i> | <i>1 to 4</i> |
| <i>UK criterion (COT)</i> | <i>2</i> |

The highest ADDs are calculated for the infants of farmer receptors and, in the absence of UK criterion for assessing exposure to dioxin-like compounds in breast milk, represent at worst less than 0.2% of the US EPA criterion of 50 pg kg⁻¹ d⁻¹ of 2,3,7,8-TCDD. The calculated ADDs for residential receptors are lower compared to the farmer since the most significant exposure to dioxins/furans is via the food chain, particularly animals and animal products. The farmer receptors are assumed to consume contaminated meat and dairy products. However, residential receptors are only assumed to consume

vegetable products which are less significant with regard to exposure to dioxins/furans.

As a worst case, the ADD for the highest exposure for the infants of farmers (Farmer Northeast) is 4.1% of the COT TDI. For these receptors it is assumed, that all of the food consumed by their mother is reared and grown locally at the location of maximum impact in their area. However, this represents an extreme worst-case. Furthermore, the duration of exposure is short and the average daily intake over the lifetime of the individual would be substantially less.

Taking into account the extreme worst-case basis for the assessment, it is concluded that infant exposure to breast milk would be not significant. Furthermore, the WHO recognises that breast-fed infants will be exposed to higher intakes for a short duration, but also that breast feeding itself provides associated benefits.

5 SUMMARY AND CONCLUSIONS

5.1 SUMMARY

The possible impacts on human health arising from dioxins and furans (PCDD/F) and dioxin-like PCBs emitted from the two SWIP at the Avonmouth site have been assessed under the worst-case scenario, namely that of an individual exposed for a lifetime to the effects of the highest airborne concentrations and consuming mostly locally grown food. This equates to a hypothetical farmer consuming food grown on the farm, situated at the closest proximity to the facility. Where there are no active farming areas in close proximity, a residential receptor is considered where it is assumed that the resident consumes locally grown vegetables.

The assessment has identified and considered the most plausible pathways of exposure for the individuals considered (farmer and resident). Deposition and subsequent uptake of the compounds of potential concern (COPCs) into the food chain is likely to be the more numerically significant pathway over direct inhalation.

The maximum combined contribution of the two SWIP to the COT TDI is 0.5% for the farmer receptors and <0.1% for the residential receptors. For the farmer this assumes as a worst-case that these receptors are located at the closest farming area to the site and all of their food is reared and grown at this location and represents an extreme worst-case. Therefore, taking into account the extreme worst-case assumptions, the impact of emissions on local sensitive receptors is considered to be not significant.

5.2 CONCLUSIONS

The risk assessment methodology used in this assessment has been structured so as to create worst case estimates of risk. A number of features in the methodology give rise to this degree of conservatism. It has been demonstrated that for the maximally exposed individual, exposure to dioxins, furans and dioxin-like PCBs is not significant.

ANNEX A

SITE PARAMETERS

Annex A: Site Parameters Defined for the Health Risk Assessment

| Parameter | Parameter Value | IRAP Symbol | Units |
|--|-----------------|------------------|--|
| Soil dry bulk density | 1.5 | bd | g cm ⁻³ |
| Forage fraction grown on contam. soil eaten by CATTLE | 1.0 | beef_fi_forage | -- |
| Grain fraction grown on contam. soil eaten by CATTLE | 1.0 | beef_fi_grain | -- |
| Silage fraction grown on contam. eaten by CATTLE | 1.0 | beef_fi_silage | -- |
| Qty of forage eaten by CATTLE each day | 8.8 | beef_qp_forage | kg DW d ⁻¹ |
| Qty of grain eaten by CATTLE each day | 0.47 | beef_qp_grain | kg DW d ⁻¹ |
| Qty of silage eaten by CATTLE each day | 2.5 | beef_qp_silage | kg DW d ⁻¹ |
| Grain fraction grown on contam. soil eaten by CHICKEN | 1.0 | chick_fi_grain | -- |
| Qty of grain eaten by CHICKEN each day | 0.2 | chick_qp_grain | kg DW d ⁻¹ |
| Fish lipid content | 0.07 | f_lipid | -- |
| Fraction of CHICKEN's diet that is soil | 0.1 | fd_chicken | -- |
| Universal gas constant | 8.205e-5 | gas_r | atm-m ³ mol ⁻¹ K ⁻¹ |
| Plant surface loss coefficient | 18 | kp | a ⁻¹ |
| Fraction of mercury emissions NOT lost to the global cycle | 0.48 | merc_q_corr | -- |
| Fraction of mercury speciated into methyl mercury in produce | 0.22 | mercmethyl_ag | -- |
| Fraction of mercury speciated into methyl mercury in soil | 0.02 | mercmethyl_sc | -- |
| Forage fraction grown contam. soil, eaten by MILK CATTLE | 1.0 | milk_fi_forage | -- |
| Grain fraction grown contam. soil, eaten by MILK CATTLE | 1.0 | milk_fi_grain | -- |
| Silage fraction grown contam. soil, eaten by MILK CATTLE | 1.0 | milk_fi_silage | -- |
| Qty of forage eaten by MILK CATTLE each day | 13.2 | milk_qp_forage | kg DW d ⁻¹ |
| Qty of grain eaten by MILK CATTLE each day | 3.0 | milk_qp_grain | kg DW d ⁻¹ |
| Qty of silage eaten by MILK CATTLE each day | 4.1 | milk_qp_silage | kg DW d ⁻¹ |
| Averaging time | 1 | milkfat_at | a |
| Body weight of infant | 9.4 | milfat_bw_infant | kg |
| Exposure duration of infant to breast milk | 1 | milkfat_ed | a |
| Proportion of ingested dioxin that is stored in fat | 0.9 | milkfat_f1 | -- |
| Proportion of mothers weight that is fat | 0.3 | milkfat_f2 | -- |
| Fraction of fat in breast milk | 0.04 | milkfat_f3 | -- |
| Fraction of ingested contaminant that is absorbed | 0.9 | milkfat_f4 | -- |
| Half-life of dioxin in adults | 2555 | milkfat_h | d |
| Ingestion rate of breast milk | 0.688 | milkfat_ir_milk | kg d ⁻¹ |
| Viscosity of air corresponding to air temp. | 1.81e-04 | mu_a | g cm ⁻¹ s ⁻¹ |
| Fraction of grain grown on contam. soil eaten by PIGS | 1.0 | pork_fi_grain | -- |
| Fraction of silage grown on contam. soil and eaten by PIGS | 1.0 | pork_fi_silage | -- |
| Qty of grain eaten by PIGS each day | 3.3 | pork_qp_grain | kg DW d ⁻¹ |
| Qty of silage eaten by PIGS each day | 1.4 | pork_qp_silage | kg DW d ⁻¹ |
| Qty of soil eaten by CATTLE | 0.5 | qs_beef | kg d ⁻¹ |
| Qty of soil eaten by CHICKEN | 0.022 | qs_chick | kg d ⁻¹ |
| Qty of soil eaten by DAIRY CATTLE | 0.4 | qs_milk | kg d ⁻¹ |
| Qty of soil eaten by PIGS | 0.37 | qs_pork | kg d ⁻¹ |
| Density of air | 1.2e-3 | rho_a | g cm ⁻³ |
| Solids particle density | 2.7 | rho_s | g cm ⁻³ |
| Interception fraction - edible portion ABOVEGROUND | 0.39 | rp | -- |
| Interception fraction - edible portion FORAGE | 0.5 | rp_forage | -- |
| Interception fraction - edible portion SILAGE | 0.46 | rp_silage | -- |
| Ambient air temperature | 298 | t | K |
| Temperature correction factor | 1.026 | theta | -- |
| Soil volumetric water content | 0.2 | theta_s | mL cm ⁻³ |
| Length of plant expos. to depos. - ABOVEGROUND | 0.16 | tp | a |
| Length of plant expos. to depos. - FORAGE | 0.12 | tp_forage | a |
| Length of plant expos. to depos. - SILAGE | 0.16 | tp_silage | a |
| Average annual wind speed | 3.9 | u | m s ⁻¹ |
| Dry deposition velocity | 0.5 | vdv | cm s ⁻¹ |
| Dry deposition velocity for mercury | 2.9 | vdv_hg | cm s ⁻¹ |
| Wind velocity | 3.9 | w | m s ⁻¹ |
| Yield/standing crop biomass - edible portion ABOVEGROUND | 2.24 | yp | kg DW m ⁻² |
| Yield/standing crop biomass - edible portion FORAGE | 0.24 | yp_forage | kg DW m ⁻² |
| Yield/standing crop biomass - edible portion SILAGE | 0.8 | yp_silage | kg DW m ⁻² |
| Soil mixing zone depth | 2.0 | z | cm |

ANNEX B

SCENARIO PARAMETERS

Annex B: Exposure Scenario Parameters

| Parameter Description | Adult Resident | Child Resident | Adult Farmer | Child Farmer | Adult Fisher | Child Fisher | Units |
|---|----------------|----------------|--------------|--------------|--------------|--------------|--|
| Averaging time for carcinogens | 70 | 70 | 70 | 70 | 70 | 70 | a |
| Averaging time for noncarcinogens | 30 | 6 | 40 | 6 | 30 | 6 | a |
| Consumption rate of BEEF | 0.0 | 0.0 | 0.00122 | 0.00075 | 0.0 | 0.0 | kg kg ⁻¹ FW d ⁻¹ |
| Body weight | 70 | 15 | 70 | 15 | 70 | 15 | kg |
| Consumption rate of POULTRY | 0.0 | 0.0 | 0.00066 | 0.00045 | 0.0 | 0.0 | kg kg ⁻¹ FW d ⁻¹ |
| Consumption rate of ABOVEGROUND PRODUCE | 0.00032 | 0.00077 | 0.00047 | 0.00113 | 0.00032 | 0.00077 | kg kg ⁻¹ DW d ⁻¹ |
| Consumption rate of BELOWGROUND PRODUCE | 0.00014 | 0.00023 | 0.00017 | 0.00028 | 0.00014 | 0.00023 | kg kg ⁻¹ DW d ⁻¹ |
| Consumption rate of DRINKING WATER | 1.4 | 0.67 | 1.4 | 0.67 | 1.4 | 0.67 | L d ⁻¹ |
| Consumption rate of PROTECTED ABOVEGROUND PRODUCE | 0.00061 | 0.0015 | 0.00064 | 0.00157 | 0.00061 | 0.0015 | kg kg ⁻¹ DW d ⁻¹ |
| Consumption rate of SOIL | 0.0001 | 0.0002 | 0.0001 | 0.0002 | 0.0001 | 0.0002 | kg d ⁻¹ |
| Exposure duration | 30 | 6 | 40 | 6 | 30 | 6 | yr |
| Exposure frequency | 350 | 350 | 350 | 350 | 350 | 350 | d a ⁻¹ |
| Consumption rate of EGGS | 0.0 | 0.0 | 0.00075 | 0.00054 | 0.0 | 0.0 | kg kg ⁻¹ FW d ⁻¹ |
| Fraction of contaminated ABOVEGROUND PRODUCE | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | -- |
| Fraction of contaminated DRINKING WATER | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | -- |
| Fraction contaminated SOIL | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | -- |
| Consumption rate of FISH | 0.0 | 0.0 | 0.0 | 0.0 | 0.00125 | 0.00088 | kg kg ⁻¹ FW d ⁻¹ |
| Fraction of contaminated FISH | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | 1.0 | -- |
| Inhalation exposure duration | 30 | 6 | 40 | 6 | 30 | 6 | a |
| Inhalation exposure frequency | 350 | 350 | 350 | 350 | 350 | 350 | d a ⁻¹ |
| Inhalation exposure time | 24 | 24 | 24 | 24 | 24 | 24 | h d ⁻¹ |
| Fraction of contaminated BEEF | 1 | 1 | 1 | 1 | 1 | 1 | -- |
| Fraction of contaminated POULTRY | 1 | 1 | 1 | 1 | 1 | 1 | -- |
| Fraction of contaminated EGGS | 1 | 1 | 1 | 1 | 1 | 1 | -- |
| Fraction of contaminated MILK | 1 | 1 | 1 | 1 | 1 | 1 | -- |
| Fraction of contaminated PORK | 1 | 1 | 1 | 1 | 1 | 1 | -- |
| Inhalation rate | 0.83 | 0.30 | 0.83 | 0.30 | 0.83 | 0.30 | m ³ h ⁻¹ |
| Consumption rate of MILK | 0.0 | 0.0 | 0.01367 | 0.02268 | 0.0 | 0.0 | kg kg ⁻¹ FW d ⁻¹ |
| Consumption rate of PORK | 0.0 | 0.0 | 0.00055 | 0.00042 | 0.0 | 0.0 | kg kg ⁻¹ FW d ⁻¹ |
| Time period at the beginning of combustion | 0 | 0 | 0 | 0 | 0 | 0 | a |
| Length of exposure duration | 30 | 6 | 40 | 6 | 30 | 6 | a |



Gair Consulting Limited
Osterley
White Lackington
Piddletrenthide
Dorchester
DT2 7QU

Tel: 01869 278889